

Stefano Goffredo · Zvy Dubinsky
Editors

The Mediterranean Sea

Its history and present challenges

Christine H.L. Schönberg and Max Wisshak

Abstract

Research conducted in the Mediterranean significantly contributed to our understanding of bioerosion, providing faunistic records and key information about the succession that occurs when fresh substrate is colonized by eroding biota. Bioeroders that have a substantial role in the Mediterranean are microendoliths, sponges, boring mollusks and various grazers. A multitude of environmental factors controls their abundances, diversities and eroding capacities. With ongoing climate change, several of these factors are likely to magnify the effects of bioerosion in the Mediterranean and worldwide. We regard eutrophication as the most important in the Mediterranean, but climate change, especially ocean acidification, will also have an important effect. Should bioerosion levels change, characteristic limestone coasts will be impacted, as will be community and sediment compositions, enigmatic cold-water coral ecosystems, mollusk aquaculture and man-made materials that are submerged. Understudied topics in Mediterranean bioerosion include rates, interactions at community level, as well as direct effects of climate change.

Keywords

Bioeroder densities • Bioeroders • Bioeroding organisms • Bioeroding taxa • Bioerosion rates of microbes • Bioerosion rates • Bioerosion • Chemical bioerosion • Coastal damage • Degradation of hard substrates • Eutrophication and pollution • Global warming • Lithic bioerosion • Macrobioerosion • Mediterranean bioerosion • Microbioerosion • Microendolithic bioerosion • Microendoliths • Ocean acidification • Sponge bioerosion • Wood borers

C.H.L. Schönberg (✉)
Australian Institute of Marine Science (AIMS),
Oceans Institute at the University of Western Australia,
Crawley, WA 9006, Australia
e-mail: C.Schoenberg@aims.gov.au

M. Wisshak
Marine Research Department, Senckenberg am Meer,
26382 Wilhelmshaven, Germany

Bioerosion in the Mediterranean

The Past of Mediterranean Bioerosion

Endolithic microbes belong to the oldest known life forms on earth with a confirmed age of at least 3.35 Ga (Fliegel et al. 2010). Endolithic cyanobacterial body fossils reach back to the Proterozoic (Golubic and Seong-Joo 1999), other microendoliths appeared in the Cambrian (Glaub and Vogel 2004). ‘Deep borers’ such as mollusks and worms were first reported from the Silurian (Beuck et al. 2008), and mobile, grazing epiliths from the Triassic (Wilson 2006). According to Bertling (1999, 2002), significant levels of bioerosion in prehistoric Europe co-occurred with the arrival of reefal substrate, and fluctuated with environmental conditions such as sedimentation levels and oxygenation, which favored or suppressed different bioeroders. Sea-level change resulted in changing successions of bioeroder communities and created a high diversity (Santos et al. 2008 and references therein). Bertling (1999, 2002) related increasing evidence of bioerosion to higher turbidity and nutrient levels, which is a recurrent theme in the past of bioerosion (e.g. Highsmith 1980a; Hallock and Schlager 1986; Hallock 1988). Being very responsive to changes in nutrients and having bathymetric distribution ranges, bioeroding sponges and their traces have probably received most attention in the Mediterranean paleontological record (Hartman 1957; Bromley and D’Alessandro 1984, 1990; Wisshak 2008). But due to their narrow ecological niche separation, microendoliths are even better paleoindicators for nutrients, temperature, light, moisture and salinity, particularly from the Late Ordovician onwards (e.g. Perry and Macdonald 2002; Vogel and Brett 2009; Wisshak 2012).

State of Art of Marine Bioerosion Research – What Is Contributed from the Mediterranean?

Since Neumann’s (1966) definition of ‘bioerosion’ as the **degradation of hard substrates by living organisms**, related research has steadily increased (Schönberg and Tapanila 2006, their Fig. 1). While bioerosion research in general is mainly concentrating on tropical coral reefs (8.5 % of the publications are from the Great Barrier Reef alone), the Mediterranean is nevertheless well represented with almost 11 % (Fig. 26.1a). In numbers of publications on bioerosion listed in Thomson Reuter’s Web of Knowledge (2011) for the last decade, study locations in southern France and Italy are on place 3 and 4 worldwide (Fig. 26.1b, however, ‘France’ data include publications on studies conducted in French Polynesia and at Reunion Island). Sites along the

Mediterranean coast of Africa were not represented, but European countries contributed about 78 % of studies conducted in the Mediterranean Sea, and Asian countries 22 % (Fig. 26.1b).

As in other areas of research, studies on bioerosion evolved from more descriptive to more applied topics, and the overall knowledge is somewhat patchy and sometimes even contradictive (Schönberg 2008). Presently, faunistic studies on Mediterranean bioeroders are well represented, especially for the Aegean Sea, but functional studies are comparatively rare. Some of the few results available indicate that grazers control settlement success in bioeroders (e.g. Bromley et al. 1990), while other workers claim that by reducing algal cover, grazers are enhancing bioeroder growth (e.g. Cebrian and Uriz 2006). Results on bioerosion rates are scarce from the Mediterranean, with one publication on sponge bioerosion (Calcinai et al. 2011) and a thesis with results on accretion and bioerosion in the Ionian Sea in relation to light and temperature with water depth (Pyko 2009). Long-term studies described the succession of bioeroders in marble blocks over time (e.g. Bromley et al. 1990), with one taxon group preparing the substrate for the next, although it has never been explained what is involved in such a preparation (see Hutchings 2008). In the Mediterranean, but also elsewhere, we do not yet fully understand why endolithic bioeroders evolved to live within hard substrates (e.g. Golubic et al. 1975; Tribollet 2008a). Proposed reasons for this life style include: protection against excessive light or water turbulence, avoidance of predators, competition, parasites and disease, saving maintenance costs, and obtaining nutrients from the substrate (Yonge 1963; Bromley 1970; Ward and Risk 1977; Vénec-Peyré 1996; Glynn 1997; Golubic et al. 2005; Tribollet 2008a; Schönberg and Wisshak 2012). The *Lithophaga* etching agent is known (a calcium-binding mucoprotein, Jaccarini et al. 1968), and microendolithic bioerosion has lately received critical attention (Garcia-Pichel 2006; Garcia-Pichel et al. 2010). However, for most bioeroder taxa using chemical bioerosion we do not know how they proceed, especially at cell level (Vénec-Peyré 1996; Martin and Britayev 1998; Golubic et al. 2005; Hutchings 2008; McLoughlin et al. 2008; Schönberg 2008). And lastly: Because of their endolithic life style, bioeroders are usually ignored in general studies and biodiversity assessments. In combination with their often difficult (ichno) taxonomy and with many species insufficiently or not yet described (e.g. Hutchings 2008; Xavier et al. 2010), we have a long way to go to understand their effects on our seas and oceans.

Agents of Mediterranean Bioerosion

Bioeroders function as a guild (Tapanila 2008) and are linked by numerous interactions, which occasionally make it




Fig. 26.1 Study locations for bioerosion research over the last 10 years (according to Thomson Reuters Web of Knowledge 2011). (a) By ocean or sea. 'Other' is comprised to 1/3 of terrestrial research, to 2/3

of research at unspecified locations. (b) By country. Denmark incl. Greenland, USA incl. Hawaii, and France incl. French Polynesia and Reunion Island

difficult to pry apart the roles of separate agents, especially in the warm-temperate biogeographic realm which appears to have a higher biodiversity of bioeroders compared to polar high or tropical low latitudes (e.g. Wisshak et al. 2011). Overall, we regard the following bioeroder groups as the most important in the Mediterranean: microendoliths, sponges, mollusks, and sea urchins. In contrast to tropical settings, fish do not play a significant role in Mediterranean bioerosion (Ballesteros 2006). Wood borers can also be key contributors, but were beyond the scope of the present publication with focus on lithic substrates. Additional details can be obtained e.g. from Yonge (1963), Bromley (1970) and Risk and MacGeachy (1978).

Microendoliths encompass the group of microbial bioeroders such as bacteria, cyanobacteria, microalgae and fungi (e.g. Golubic et al. 1975, 2005; Risk and MacGeachy 1978; Tribollet 2008a). Cyanobacteria and chlorophyte algae are key agents (Fig. 26.2) that were investigated in detail in the Mediterranean near Marseille by Le Campion-Alsumard (1975, 1979), building upon the historic studies carried out by Bornet and Flahault (1888, 1889) and Ercegović (1927, 1929, 1932). Wherever fresh substrate becomes available it is colonized and attacked within days to weeks by pioneering microbes, which then go through a succession of different taxa, reaching an equilibrium after months to years (e.g. Hong 1980; Le Campion-Alsumard et al. 1995; Sartoretto 1998;

Naylor and Viles 2002; Beuck et al. 2010). In shallow water, substrate surfaces are grazed by mollusks and echinoids as soon as microendoliths have taken hold, intensifying the effect imparted by the microbes and shaping limestone coasts (e.g. Schneider 1976, 1977; Bromley et al. 1990). Bathymetric patterns and penetration depths within substrates can be related to the compensation depth of photosynthetic microendoliths, with only fungal and bacterial microbioerosion in aphotic depths (e.g. Golubic et al. 1975; Sartoretto 1998; Tribollet 2008b). On the other hand, where light suddenly increases it can cause blooms in the microendolithic communities, possibly producing excess nutrients passed on to neighboring or host organisms (Fine and Loya 2002; Fine et al. 2004). Pace and diversity in development and linked bioerosion rates of microbes are thus highly dependent on bathymetry and the extent of photic zones, as well as the orientation of the substrate (Wisshak et al. 2011).

Sponges are among the best studied bioeroders in the Mediterranean, especially from Spain, France and Italy (e.g. Coll et al. 2010). Excavating sponges establish in newly available substrates after about 2 years and often thrive better on the undersides of substrates or ledges, where competition with coralline algae is lower (Bromley et al. 1990; Cerrano et al. 2001). They etch fine, cup-shaped grooves and mechanically remove lentil-shaped chips that are expelled with the exhalant water to form fine-grained sediment




Fig. 26.2 SEM images of epoxy resin casts of microborings produced by the cyanobacterium *Mastigocoleus testarum* (a) and the chlorophyte alga *Ostreobium queketti* (b) in 15 and 50 m water depth, respectively,

recorded on a settlement experiment off the southern Peloponnes, Ionian Sea (Images courtesy of I. Pyko)




Fig. 26.3 The most common and important bioeroding sponges of the Mediterranean: *Cliona viridis* (a, grayish tissue) and *Cliona celata* (b, yellow tissue) (Photographs courtesy of T. Perez)

(Fütterer 1974; Rützler and Rieger 1973). The proportion of chemical etching compared to what is removed as chip has been estimated between 2:98 and 70:30 %, a discrepancy that may well be related to environmental conditions (Rützler and Rieger 1973; Zundeleovich et al. 2007; Nava and Carballo 2008). In the Mediterranean *Cliona viridis* is the most abundant and destructive sponge, partly rivaled by species of the *Cliona celata* species complex (Fig. 26.3; e.g. Rosell et al. 1999; Calcinaï et al. 2011). Bioeroding sponges have been recognized to be good pollution indicators and biomonitors for environmental conditions (e.g. Hong 1980; Carballo et al. 1994, 1996).

Many mollusks such as chitons and limpets produce ‘home scars’ or resting scars at places where they most often sit (Fig. 26.4a), and generate grazing traces by

removing substrate layer by layer either by radular scraping or by use of acid (e.g. Hutchings 1986). Their intertidal activity is strongly governed by access to water, and areas closer to the water line and spray zone are usually more strongly eroded, which can lead to coastal notch-formation (e.g. Palmer et al. 2003; Neumann 1966). Locally more important than these surface marks are the large hollows mostly produced by bivalves such as pholads, gastrochaenids and mytilids. These mollusks erode by mechanical or chemical means (e.g. Jaccarini et al. 1968; Ansell and Balakrishna Nair 1969; Appukuttan 1969). For reefal areas pholad densities between 50 and 500 individuals per m² can occur (Warne 1975; Highsmith 1980b; Loya 1982), but we have no comparable data for the Mediterranean. Here, the slow-growing mytilid lithophags play a considerable role in




Fig. 26.4 (a) *Patella caerulea* and *Chiton olivaceus* with *Patella* resting scars at the Costa Daurada, Spain. (b) Borehole battery made by *Lithophaga lithophaga* in the French Mediterranean (a – courtesy of J. Martinell, b – of A. Antonioli)




Fig. 26.5 Bioerosion caused by regular sea urchins. (a) Star-shaped scrape marks made by *Paracentrotus lividus* on an *Anomia ephippium* valve from the Costa Brava, Spain. (b) Resting scars produced by *Paracentrotus lividus* at the Costa Daurada, Spain (Photographs courtesy of J. Martinell)

weakening coastal structures (Fig. 26.4b; Kleemann 1973; Devescovi 2009), a process that is magnified by lithophags being harvested as a local delicacy (Fanelli et al. 1994; Russo and Cicogna 1991; Hrs-Brenko et al. 1991; Frascchetti et al. 2001; Guidetti et al. 2003; Devescovi et al. 2005). Therefore, despite being abundant, since 2005 *Lithophaga lithophaga* is a CITES-protected species (Boudouresque et al. 1991; CITES 2011), a restriction which is not necessarily respected by the people (Bianchi and Morri 2000).

Sea urchins appear to play a comparatively small role in Mediterranean bioerosion (Fig. 26.5; Laubier 1966; Sartoretto and Francour 1997), which may in part be related to the fact that they are locally overfished for their roe (Guidetti et al. 2004; Micael et al. 2009). But where urchins have large diameters or their densities reach high levels of at least 20 individuals per 25 m² (Sartoretto and Francour 1997)

their continuous grazing can maintain ‘barren grounds’ of mostly coralline algae (Guidetti et al. 2003; Privitera et al. 2005), they can create cup-shaped borings or scrape deep channels into the substrate surface (e.g. Asgaard and Bromley 2008; Fig. 26.5b). Where they graze, sea urchins remove live tissue of other invertebrates or infaunal bioeroders (e.g. Tribollet and Golubic 2005) and can dislodge smaller sessile fauna by bulldozing (Schönberg, pers. obs.). By their rasping activity they produce substrate debris that is defecated as sediments and will accumulate in the area, a process that has been studied in the neighboring Red Sea (Mokady et al. 1996).

Bioeroding worms should be mentioned as well, as they cause the mud blister syndrome in mollusk shells (Fig. 26.6a; e.g. Martin and Britayev 1998). Worms establish themselves in fresh substrate at a rate similar to sponges and appear in a succession of taxa after 2–3 years (e.g. Hutchings 2008).




Fig. 26.6 (a) Mud blister in an *Anomia ephippium* valve from the Costa Brava, Spain. (b) This benthic community of bioeroders in the French Mediterranean is a result of a complicated interplay of e.g.

nutrient availability to the filter feeders (*Cliona celata* and endolithic molluscs), competition with coralline algae and other benthic organisms and predation (a – courtesy of J. Martinell, b – of T. Perez)

Their occurrences vary with environmental conditions, and they can be quite useful as pollution indicators (Hong 1980; Dean 2008). New species can become introduced pests and invasive species (Simboura and Zenetos 2005; Çinar et al. 2011; Zenetos et al. 2011). Polychaetes are a comparatively prominent worm group in the Mediterranean (Feldmann 1937; Hong 1980), while only three species of local endolithic sipunculids are presently identified (Laubier 1966; Sartoretto and Francour 1997). Ironically, Mediterranean worm reefs built by other polychaetes can become subject to bioerosion (Porrás et al. 1996; Fornós et al. 1998).

Environmental Factors Acting on Mediterranean Bioerosion and Possible Trends

Many bioeroding organisms such as microendoliths, sponges, worms and endolithic mollusks are sessile and depend on nutrients and/or particulate organic matter being carried to them in the water column. But even mobile grazers can benefit from fertilization if it induces increased algal growth (e.g. Le Bris et al. 1998; Carreiro-Silva et al. 2005, 2009). We therefore assume that in the Mediterranean the strongest local drivers of future trends of bioerosion will be eutrophication and pollution (e.g. Coll et al. 2010). Both factors are thought to aggravate the effects of climate change (Le Grand and Fabricius 2011), which will be of importance in a region that is highly industrialized, densely populated, and has comparatively little exchange with surrounding water masses (e.g. Bianchi and Morri 2000). In particular urban development

and river runoff bring about anthropogenic stress known to affect shifts in community structures and to increase bioeroder densities and bioerosion rates (Le Grand and Fabricius 2011 and their table 1). However, the magnitude of nutrient effects are probably the most difficult to determine at community level, because a multitude of cascades and feed-back loops exist that may either enhance the overall effect or confuse the results (e.g.: pollution → increased bacteria concentrations in water column → increased abundances of filter feeding bioeroders → intensive recycling of nutrients → fertilization of microendoliths → increased porosity of substrate → increased efficiency of grazer bioerosion → creation of fresh surfaces aiding settlement of bioeroders; e.g. Corredor et al. 1988; Rose and Risk 1985; Carreiro-Silva et al. 2009; Fig. 26.6b). Further complications arise from the complex interplay of dissolved inorganic nutrients, particulate organic matter, siliciclastic material or pollutants such as heavy metals in the water, with different effects on the various types of bioeroding taxa (fertilization, shading, poisoning, etc.). It is thus hardly a surprise that studies on the influence of the trophic regime on bioerosion yielded partly contradictory results (e.g. Chazottes et al. 2002; Szmant 2002; Carreiro-Silva et al. 2005; Holmes et al. 2009).

To date not many data are available on how climate change may affect bioerosion and whether global warming or ocean acidification may bring about a pertinent change. Within physiological tolerance levels chemical bioerosion itself may be enhanced by increased temperatures due to the simple fact that the reaction rate is accelerated. Overall, global warming is expected to have less severe effects on bioeroding organisms than on calcifiers, because grazers will

move into shade and endoliths are extremely well sheltered and insulated by living within porous, water-retaining substrates that provide shade, and many endoliths can produce cooling currents passing through or over their bodies. Some very common species of bioeroding sponges live in symbiosis with dinoflagellates similar to those in corals, yet they have been observed to be very bleaching resistant during periods of abnormal heating (Vicente 1990; Schönberg and Wilkinson 2001), but the symbiosis may be quite susceptible to light stress (Hill and Wilcox 1998; Schönberg and Suwa 2007; Schönberg et al. 2008). If other bioeroders are as heat resistant as these sponges, the entire guild may indirectly benefit from thermal damage to the calcifiers, which in response to heat stress may be weakened or dead and thus an easy target. In the Mediterranean, warming events are expected to increase, and eastern parts of the basin are more likely to be affected than central and western parts (e.g. Bianchi and Morri 2000; Garrabou et al. 2001, 2009; Coll et al. 2010).

Ocean acidification may have a larger impact on bioerosion than global warming. Most macrobioeroders are either filter feeders or grazers, groups that were found to not be negatively affected or even more common near Mediterranean CO₂ vents (Kroeker et al. 2011). Moreover, many bioeroders at least in part use chemical dissolution to remove substrate, a process which is often thought to be pH dependent or relying on acid production (e.g. Pomponi 1980). Where this is the case, bioerosion is likely to increase in acidified waters (Tribollet et al. 2006). Experimental evidence on how bioerosion might change as a direct reaction to climate change and ocean acidification is scarce. With higher pCO₂ Tribollet et al. (2006, 2009) observed increased penetration depth and bioerosion by microendoliths with a stable community composition. Studies by the present authors investigate the impacts of climate change on macrobioerosion, and results suggest that temperature has little effect on sponge bioerosion (Wisshak et al. unpubl. data), but it will be significantly accelerated by ocean acidification (Wisshak et al. 2012). However, if acidification ever reaches catastrophic levels, reducing the abundance of suitable substrate in the Mediterranean (e.g. Fine and Tchernov 2007; Kroeker et al. 2011), lithic bioerosion will likewise come under pressure. Volcanic activity and hydrothermal seepage in the Mediterranean provides an ideal environment to study consequences of ocean acidification on bioerosion in natural settings (Kroeker et al. 2011; Rodolfo-Metalpa et al. 2011).

Which Assets Will Be Most Likely Affected by Mediterranean Bioerosion?

Mediterranean bioeroder communities are constantly modifying and structuring biogenic as well as abiogenic hardground

of coastal areas. This includes the characteristic coralline alga assemblages that are mostly shaped by grazers such as sea urchins (Fig. 26.7; Ballesteros 2006). In extreme cases the urchins can create and maintain 'barrens' devoid of high structural complexity (Privitera et al. 2005). Cover with red algae will in turn reduce settlement success of the larvae of many bioeroders and withstands bioerosion better than other substrates (Hong 1980; Smyth 1989; Chazottes et al. 2002; Ballesteros 2006), although bioeroding sponges frequently penetrate it (Cotte 1914; Cerrano et al. 2001).

Many bioeroders rework and produce considerable amounts of sediment. For example, sponge-generated sediments in the North Adriatic Sea make up about 2–3 % (Fütterer 1974). Higher bioeroder densities or shifts in their community compositions would mean that both their influence on the hard-bottom morphology and the sediment quality in their habitat will change (Hutchings 1986).

Diverse bioeroder communities inhabit slow-growing solitary, (pseudo)colonial and cold-water Mediterranean corals (Fig. 26.8a; Bromley 2005; Wisshak 2008; Mastrototaro et al. 2010), which they can infest to a high degree (20–75 %, Bavestrello et al. 1997; Corriero et al. 1997; Cerrano et al. 2001), deform them (Beuck et al. 2007), cause a higher risk of fragmentation by removing up to 70 % of the skeleton (Beuck et al. 2010), overwhelm early life stages (Calcinai et al. 2000) or cause wide-spread mortality (Corriero et al. 1997). Bioeroding sponges in particular are a devastating pest in Mediterranean corals and among others attack the precious red coral *Corallium rubrum* (Fig. 26.8b; e.g. Melone 1965; Maldonado 1992; Bavestrello et al. 1996; Calcinai et al. 2000, 2002, 2008), the overexploited Mediterranean 'red gold'. *Corallium* has been used as a gemstone since ancient times (Pronzato 2000), historically supported trade between Mediterranean countries and India, and is part of the Greek mythology (Medusa's head, Poseidon's Palace; De Simone 2010 and other contributions in the same volume). Bioeroding sponges are very common in Mediterranean *Corallium* colonies (Barletta and Vighi 1968). Corriero et al. (1997) stated that coral survival is strongly correlated with severity of infestation with these borers that are able to kill colonies that reach 4 years of age, while they only become reproductive at 2 years or later (Santangelo et al. 2003). Should the infestation rate by sponges increase, the traditional Mediterranean culture and trade based on the already threatened red precious coral may well become a thing of the past (Weinberg 1991; Garrabou et al. 2001, 2009; Costantini et al. 2007). As many cold-water reefs support a very diverse associated fauna, a reduced abundance of such corals would affect other taxa as well (e.g. Mastrototaro et al. 2010).

Bioeroders furthermore invade mollusk shells and can cause considerable damage on local oysters and mussels. Aquaculture hatcheries generate considerable profit for at least six Mediterranean countries (Fisheries and Aquaculture




Fig. 26.7 The 'coralligène' (a) and some of the main players involved in shaping this habitat (b): the sea urchin *Paracentrotus lividus*, the sponge *Cliona viridis* (brown papillae in the lower left) and coralline algae (in the background) (a – courtesy of T. Perez)




Fig. 26.8 (a) The cold-water corals *Madropora oculata* and the solitary *Desmophyllum dianthus* from about 400 m water depth in the Lacaze-Duthiers Canyon near Banyuls-sur-Mer, France. (b) Colony of the red coral *Corallium rubrum* from the Linosa Trough in the Pelagian Island

group showing clear evidence of bioerosion and subsequent deformation (a – courtesy of Senckenberg/MARUM, b – courtesy of L. Beuck and A. Freiwald)

Department 2011). France is a large, traditional oyster producer, ranking among the top producer and consumer countries of the world, with an annual production of 9500 t in the Thau Lagoon (Héral 1989; Buestel et al. 2009). Losses related to bioerosion have been historically observed (Carazzi 1895) and may significantly increase in the future. While oysters are readily attacked by various bioeroders (Labura and Hrs-Brenko 1990: up to 89 % oysters by *Polydora hoplura*, Rosell et al. 1999: 100 % of oysters by two species of sponges), mussels are to a much lesser extent and degree, but often with more devastating effects to their shell stability and thus to survival from predation (Lauckner 1983).

Bioerosion in combination with changing weather patterns and increased frequencies of severe storms may also result in

more coastal damage, because bioerosion reduces the security of attachments and breaking strength of the invaded organisms and materials (Tunncliffe 1979; Highsmith 1980b; Mitchell-Tapping 1983; Clark and Morton 1999; Scott and Risk 1988). Microendoliths and urchins can attack volcanic materials such as dolerites, balsanites and volcanic glass (Allouc et al. 1996; McLoughlin et al. 2008), which would play a role in those parts of the Mediterranean that have a geological history of volcanism (e.g. Aiello et al. 2001). And as final reminder of cause-and-response to our actions: Man-made environmental changes influencing bioerosion will also damage man-made or man-shaped substrates. A number of bioeroding organisms such as sponges and pholad bivalves are able to riddle and whittle away artificial materials or



Fig. 26.9 Greek statues from the Athens National Museum of Archaeology retrieved from a shipwreck in the Aegean. They were exposed to marine bioerosion for about 2100 years, the degree of damage depending on the sediment cover. (a) The statue's

eyes were replaced by a brachiopod and a pholad boring. (b) Parts of the statue that were not covered by sand were strongly affected by bioerosion (Photographs courtesy of R. G. Bromley and U. Asgaard)

structures (Scott 1991; Brusco et al. 2005). Other mollusks are destabilizing breakwater boulders in the Adriatic (Devescovi and Iveša 2008) and can easily penetrate concrete containing calcium, as can some sponges and polychaetes (Yonge 1963; Scott et al. 1988). Echinoids have been suspected to attack even steel pilings (Emery 1960) and are known to pry grains out of granite at a faster rate than they would erode limestone (Bromley 1970). And in conclusion, bioerosion also affects submerged works of art, e.g. Greek marble statues or submerged mosaics in the Aegean (Fig. 26.9; Bromley and Asgaard 2004; Ricci et al. 2008).

Acknowledgements The authors thank Z. Dubinsky and S. Goffredo for inviting us to contribute a chapter on bioerosion to this book, and L. Walker for unearthing some of the literature on Mediterranean mud blister worms. We are indebted to the following colleagues who gave us permission to use some of their photographs: A. Antonioli, L. Beuck and A. Freiwald (IPAL 2006 and ROV Quest, MARUM 2006), T. Perez, and I. Pyko, J. Martinell, and R. Bromley and U. Asgaard.

References

- Aiello G, Budillon V, Cristofalo G, D'Argenio B, de Alteriis G, De Lauro M, Ferraro L, Marsella E, Pelosi N, Sacchi M, Tomielli R (2001) Marine geology and morphobathymetry in the Bay of Naples (south-eastern Tyrrhenian Sea, Italy). In: Faranda FM, Guglielmo L, Spezie G (eds) Mediterranean ecosystems. Structures and processes. Springer, Milano, pp 1–8
- Allouc J, Le Campion Alsumard T, Tack DL (1996) Bioerosion of magmatic rock in a coastal environment: the example of the Cap Vert Peninsula (western Senegal). *Geobios* 29:485–502
- Ansell AD, Balakrishna Nair N (1969) A comparative study of bivalves which bore mainly by mechanical means. *Am Zool* 9:857–868
- Appukuttan KK (1969) Coral-boring bivalves of Gulf of Mannar and Palk Bay. In: Proceedings of the symposium corals and coral reefs India, Marine Biological Association of India, Mandapam Camp, India, pp 379–398
- Asgaard U, Bromley RG (2008) Echinometrid sea urchins, their trophic styles and corresponding bioerosion. In: Wisshak M, Tapanila L (eds) Current developments in bioerosion. Springer, Heidelberg, pp 279–303
- Ballesteros E (2006) Mediterranean coralligenous assemblages: a synthesis of present knowledge. *Oceanogr Mar Biol Ann Rev* 44:123–195
- Barletta G, Vighi M (1968) Recherche sul corallo rosso: V. Poriferi perforanti lo sclerasse di *Corallium rubrum* Lamarck. *Rend Ist Lombardo Sci Lett Milano B* 102:145–159
- Bavestrello G, Calcinai B, Sarà M (1996) *Delectona ciconiae* sp. nov. (Porifera, Demospongiae) boring in the scleraxis of *Corallium rubrum*. *J Mar Biol Assoc UK* 76:867–873
- Bavestrello G, Calcinai B, Cerrano C, Sarà M (1997) *Delectona madreporica* n. sp. (Porifera, Demospongiae) boring the corallites of some scleractinians from the ligurian Sea. *Ital J Zool* 64:273–277
- Bertling M (1999) Late Jurassic reef bioerosion – the drawing of a new era. *Bull Geol Soc Den* 45:173–176
- Bertling M (2002) Coral reef bioerosion in times of crises – the Late Triassic/Early Jurassic example. In: Proceedings of the 9th international coral reef symposium, Bali, vol 1, pp 23–27
- Beuck L, Vertino A, Stepina E, Karolczak M, Pfannkuche O (2007) Skeletal response of *Lophelia pertusa* (Scleractinia) to bioeroding sponge infestation visualised with micro-computed tomography. *Facies* 53:157–176
- Beuck L, Wisshak M, Munnecke A, Freiwald A (2008) A giant boring in a Silurian stromatoporoid analysed by computer tomography. *Acta Palaeontol Pol* 53:147–158
- Beuck L, Freiwald A, Taviani M (2010) Spatiotemporal bioerosion patterns in deep-water scleractinians from off Santa Maria die Leuca (Apulia, Ionian Sea). *Deep-Sea Res II* 57:458–470
- Bianchi CN, Morri C (2000) Marine biodiversity of the Mediterranean Sea: situation, problems and prospects for future research. *Mar Poll Bull* 40:367–376
- Bornet E, Flahault C (1888) Note sur deux nouveaux genres d'algues perforantes. *J Bot Morot* 2:161–165
- Bornet E, Flahault C (1889) Sur quelques plantes vivant dans le test calcaire des mollusques. *Bull Soc Bot France* 36:147–179
- Boudouresque CF, Avon M, Gravez V (1991) Les espèces marines à protéger en Méditerranée. GIS Posidonie, Marseille, pp 1–447

- Bromley RG (1970) Borings as trace fossils and *Entobia cretacea* Portlock, as an example. In: Crimes TP, Harper JC (eds) Trace fossils. Geol J special issue 3:49–90
- Bromley RG (2005) Preliminary study of bioerosion in the deep-water coral *Lophelia*, Pleistocene, Rhodes, Greece. In: Freiwald A, Roberts JM (eds) Cold-water corals and ecosystems. Springer, Heidelberg, pp 898–914
- Bromley RG, Asgaard, U (2004) Archaeological disaster a bioerosion goldmine: test-blocks retrieved 2100 years after deposition, Antikythira, Greece. In: Mikuláš R (ed) Abstracts for the 4th bioerosion workshop, Czech Republic, pp 5–6
- Bromley RG, D'Alessandro A (1984) The ichnogenus *Entobia* from the Miocene, Pliocene and Pleistocene of southern Italy. Riv It Paleont Stratigr 90:227–296
- Bromley RG, D'Alessandro A (1990) Comparative analysis of bioerosion in deep and shallow water, Pliocene to recent, Mediterranean Sea. Ichnos 1:43–49
- Bromley RG, Hanken N-M, Asgaard U (1990) Shallow marine bioerosion: preliminary results of an experimental study. Bull Geol Soc Den 38:85–99
- Brusco F, Calcinai B, Bavestrello G, Cerrano C (2005) Attività di erosione di *Cliona celata* Grant, 1826 e *Pione vastifica* (Hancock, 1849) (Porifera, Demospongiae) perforanti substrati artificiali. Biol Mar Medit 12:240–243
- Buestel D, Ropert M, Prou J, Goulletquer P (2009) History, status, and future of oyster culture in France. J Shellfish Res 28:813–820
- Calcinai B, Cerrano C, Bavestrello G, Milanese M, Sarà M (2000) Il popolamento di spugne perforatrici di *Corallium rubrum* e di alcuni madreporari del Promontorio di Portofino. Boll Mus Inst Biol Univ Genova 64–65:53–59
- Calcinai B, Cerrano C, Bavestrello G (2002) A new species of *Scantiletta* (Demospongiae, Clionidae) from the Mediterranean precious coral with some remarks on the genus. Bull Mar Sci 70:919–926
- Calcinai B, Cerrano C, Iwasaki N, Bavestrello G (2008) Sponges boring into precious corals: an overview with description of a new species of *Alectona* (Demospongiae, Alectonidae) and a worldwide identification key for the genus. Mar Ecol 29:273–279
- Calcinai B, Bavestrello G, Cuttone G, Cerrano C (2011) Excavating sponges from the Adriatic Sea: description of *Cliona adriatica* sp. nov. (Demospongiae: Clionidae) and estimation of its boring activity. J Mar Biol Assoc UK 91:339–346
- Carazzi D (1895) Revisione del genere *Polydora* Bosc e cenni su due specie che vivono sulle ostriche. Mitt zool Stat Neapel 11:4–45
- Carballo JL, Sanchez-Moyano JE, García-Gómez JC (1994) Taxonomic and ecological remarks on boring sponges (Clionidae) from the Straits of Gibraltar (southern Spain): tentative bioindicators? Zool J Linn Soc Lond 112:407–424
- Carballo JL, Naranjo SA, García-Gómez JC (1996) Use of marine sponges as stress indicators in marine ecosystems at Algeciras Bay (southern Iberian Peninsula). Mar Ecol Prog Ser 135:109–122
- Carreiro-Silva M, McClanahan TR, Kiene WE (2005) The role of inorganic nutrients and herbivory in controlling microbioerosion of carbonate substratum. Coral Reefs 24:214–221
- Carreiro-Silva M, McClanahan TR, Kiene WE (2009) Effects of inorganic nutrients and organic matter on microbial euendolithic community composition and microbioerosion rates. Mar Ecol Prog Ser 392:1–15
- Cebrian E, Uriz M-J (2006) A facilitation example involving sponges: grazing on fleshy seaweeds by sea urchins enhances *Cliona viridis* abundance. Mar Ecol Prog Ser 323:83–89
- Cerrano C, Bavestrello G, Bianchi CN, Calcinai B, Cattaneo-Vietti R, Morri C, Sarà M (2001) The role of sponge bioerosion in Mediterranean coralligenous accretion. In: Faranda FM, Guglielmo L, Spezie G (eds) Mediterranean ecosystems. Structures and processes. Springer, Milano, pp 235–240
- Chazottes V, Le Campion-Alsumard T, Peyrot-Clausade M, Cuet P (2002) The effects of eutrophication-related alterations to coral reef communities on agents and rates of bioerosion (Reunion Island, Indian Ocean). Coral Reefs 21:375–390
- Çinar ME, Bilecenoglu M, Öztürk B, Kayağan T, Yokeş MB, Aysel V, Dağlı E, Açık S, Özcan T, Erdoğan H (2011) An updated review of alien species on the coasts of Turkey. Medit Mar Sci 12:257–315
- CITES (2011) Consideration of proposals for amendment of Appendices I and II, and Appendices I, II and II valid from 27 April 2011. <http://www.cites.org/eng/cop/13/prop/e13-p35.pdf>; and <http://www.cites.org/eng/app/E-Apr27.pdf>. Accessed 14 Sept 2011
- Clark T, Morton B (1999) Relative role of bioerosion and typhoon-induced disturbance on the dynamics of a high latitude scleractinian coral community. J Mar Biol Assoc UK 79:83–820
- Coll M, Piroddi C, Steenbeek J, Kaschner K, Ben Rais Lasram F, Aguzzi J, Ballesteros E, Bianchi CN, Corbera J, Dailianis T, Danovaro R, Estrada M, Froglio B, Galil BS, Gasol JM, Gertwagen R, Gil J, Guilhaumon F, Kesner-Reyes K, Kitsos M-S, Koukouras A, Lampadariou N, Laxamana E, de la Cuadra L-F, Lotze HK, Martin D, Mouillot D, Oro D, Raicevich S, Ruis-Barile J, Saiz-Salinas JJ, San Vicente C, Somot S, Templado J, Turon X, Vafidis D, Villanueva R, Voultsiadou E (2010) The biodiversity of the Mediterranean Sea: estimates, patterns, and threats. PLoS One 5(e11842):1–36
- Corredor JE, Wilkinson CR, Vicente VP, Morell JM, Otero E (1988) Nitrate release by Caribbean reef sponges. Limnol Oceanogr 33:114–120
- Corriero G, Abbiati M, Santangelo G (1997) Sponges inhabiting a Mediterranean red coral population. PSZN Mar Ecol 18:147–155
- Costantini F, Fauvelot C, Abbiati M (2007) Genetic structuring of the temperate gorgonian coral (*Corallium rubrum*) across the Western Mediterranean Sea revealed by microsatellites and nuclear sequences. Mol Ecol 16:5168–5182
- Cotte J (1914) L'association de *Cliona viridis* (Schmidt) (spong.) et de *Lithophyllum expansum* (Philippi) (algues). Comptes Rendus des Séances de la Société Biologie Paris 76:739–740
- De Simone AE (2010) 'The Coral Routes': From the Mediterranean Sea to the world-wide jewelry. In: Bussoletti E, Cottingham D, Bruckner A, Roberts G, Sandulli R (eds) Proceedings of the international workshop on red coral science, management and trade. Lessons from the Mediterranean Naples 2009, NOAA Technical Memorandum CRCP-13, Silver Spring, MD, USA, pp 102–105
- Dean HK (2008) The use of polychaetes (Annelida) as indicator species of marine pollution: a review. Rev Biol Trop 56:11–38
- Devescovi M (2009) Biometric differences between date mussels *Lithophaga lithophaga* colonizing artificial and natural structures. Acta Adriatica 50:129–138
- Devescovi M, Iveša L (2008) Colonization patterns of the date mussel *Lithophaga lithophaga* (L., 1758) on limestone breakwater boulders of a marina. Periodicum Biol 110:339–345
- Devescovi M, Ozretić B, Iveša L (2005) Impact of date mussel harvesting on the rocky bottom structural complexity along the Istrian coast (northern Adriatic, Croatia). J Exp Mar Biol Ecol 325:134–145
- Emery KO (1960) The sea off southern California, a modern habitat of petroleum. Wiley, New York, p 147
- Ercegović A (1927) Tri nova roda litofitskih cijanoficeja sa jadranske obale. Acta Bot Zagreb Univ 2:78–84
- Ercegović A (1929) Sur quelques nouveaux types des Cyanophycées lithophytes de la côte Adriatique. Arch Protistenkunde 66:164–174
- Ercegović A (1932) Ekološke i sociološke studije o litofitskim cijanoficejama sa jugoslavenske obale Jadrana. Rad Jugoslav akad znan umjet. Mat-prirod razred Knj 75:129–220
- Fanelli G, Piraino S, Belmonte G, Geraci S, Boero F (1994) Human predation along Apulian rocky coast: desertification caused by *Lithophaga lithophaga* (Mollusca) fisheries. Mar Ecol Prog Ser 110:1–8
- Feldmann J (1937) Recherches sur la végétation marine de la Méditerranée: la côte des Albères. Wolf, Rouen, p 339

- Fine M, Loya Y (2002) Endolithic algae: an alternative source of photoassimilates during coral bleaching. *Proc R Soc Lond* 269:1205–1210
- Fine M, Tchernov D (2007) Scleractinian coral species survive and recover from decalcification. *Science* 315:1811
- Fine M, Steindler L, Loya L (2004) Endolithic algae photoacclimate to increased irradiance during bleaching. *Mar Freshw Res* 55:115–121
- Fisheries and Aquaculture Department (2011) Mediterranean fisheries. <http://www.fao.org/docrep/006/N7865E/N7865E01.htm>. Accessed 13 July 2011
- Fliegel D, Kosler J, McLoughlin N, Simonetti A, de Wit MJ, Wirth R, Furnes H (2010) In-situ dating of the earth's oldest trace fossil at 3.34 Ga. *Earth Planet Sci Lett* 299:290–298
- Fornós JJ, Forteza V, Martínez-Taberner A (1998) Modern polychaete reefs in Western Mediterranean lagoons: *Ficopotamus enigmatus* (Fauvel) in the Albufera of Menorca, Balearic Islands. *Palaeogeogr Palaeoclimatol Palaeoecol* 128:175–186
- Fraschetti S, Bianchi CN, Terlizzi A, Fanelli G, Morri C, Boero F (2001) Spatial variability and human disturbance in shallow subtidal hard substrate assemblages: a regional approach. *Mar Ecol Prog Ser* 212:1–12
- Fütterer DK (1974) Significance of the boring sponge *Cliona* for the origin of fine grained material of carbonate sediments. *J Sed Petrol* 48:1149–1157
- Garcia-Pichel F (2006) Plausible mechanisms for the boring in carbonates by microbial phototrophs. *Sed Geol* 185:205–213
- Garcia-Pichel F, Ramirez-Reinat E, Gao Q (2010) Microbial excavation of solid carbonates powered by P-type ATPase-mediated transcellular Ca^{2+} transport. *PNAS* 107:21749–21754
- Garrahou J, Pérez T, Sartoretto S, Harmelin JG (2001) Mass mortality event in red coral *Corallium rubrum* populations in the Provence region (France, Mediterranean). *Mar Ecol Prog Ser* 217:263–272
- Garrahou J, Coma R, Bensoussan N, Bally M, Chevaldonné P, Cigliano M, Diaz D, Harmelin JG, Gambi MC, Kersting DK, Ledoux JB, LeJeune C, Linares C, Marschall C, Pérez T, Ribes M, Romano JC, Serrano E, Teixido N, Torrents O, Zabala M, Zuberer F, Cerrano C (2009) Mass mortality in northwestern Mediterranean rocky benthic communities: effects of the 2003 heat wave. *Glob Chang Biol* 15:1090–1103
- Glaub I, Vogel K (2004) The stratigraphic record of microborings. *Fossils Strata* 51:126–135
- Glynn PW (1997) Bioerosion and coral reef growth: a dynamic balance. In: Birkeland C (ed) *Life and death of coral reefs*. Chapman and Hall, New York, pp 69–98
- Golubic S, Seong-Joo L (1999) Early cyanobacterial fossil record: preservation, palaeoenvironments and identification. *Eur J Phycol* 34:339–348
- Golubic S, Perkins RD, Lukas KJ (1975) Boring microorganisms and microborings in carbonate substrates. In: Frey EW (ed) *The study of trace fossils*. Springer, New York, pp 229–259
- Golubic S, Radtke G, Le Campion-Alsumard T (2005) Endolithic fungi in marine ecosystems. *Trends Microbiol* 13:229–235
- Guidetti P, Fraschetti S, Terlizzi A, Boero F (2003) Distribution patterns of sea urchins and barrens in shallow Mediterranean rocky reefs impacted by the illegal fishery of the rock-boring mollusc *Lithophaga lithophaga*. *Mar Biol* 143:1135–1142
- Guidetti P, Terlizzi A, Boero F (2004) Effects of the edible sea urchin, *Paracentrotus lividus*, fishery along the Apulian rocky coast (SE Italy, Mediterranean Sea). *Fish Res* 66:287–297
- Hallock P (1988) The role of nutrient availability in bioerosion: consequences to carbonate buildups. *Palaeogeogr Palaeoclimatol Palaeoecol* 63:275–291
- Hallock P, Schlager W (1986) Nutrient excess and the demise of coral reefs and carbonate platforms. *Palaios* 1:389–398
- Hartman W (1957) Ecological niche differentiation in the boring sponges (Clionidae). *Evolution* 11:294–297
- Héral M (1989) Traditional oyster culture in France. In: Barnabé G, Solbé JF, Laird L (eds) *Aquaculture*, vol 1. Ellis Horwood, London, pp 342–387
- Highsmith RC (1980a) Geographic patterns of coral bioerosion: a productivity hypothesis. *J Exp Mar Biol Ecol* 46:177–196
- Highsmith RC (1980b) Passive colonization and asexual colony multiplication in the massive coral *Porites lutea* Milne Edwards & Haime. *J Exp Mar Biol Ecol* 47:55–67
- Hill M, Wilcox T (1998) Unusual mode of symbiont repopulation after bleaching in *Anthosigmella varians*: acquisition of different zooxanthellae strains. *Symbiosis* 25:279–289
- Holmes G, Ortiz J-C, Schönberg CHL (2009) Bioerosion rates of the sponge *Cliona orientalis* Thiele, 1900: spatial variation over short distances. *Facies* 55:203–211
- Hong JS (1980) Étude faunistique d'un fond de concrétionnement de type coralligène soumis à un gradient de pollution en Méditerranée nord-occidentale (Golfe de Fos). Dissertation, University Aix-Marseille II, 134 pp
- Hrs-Brenko M, Zavodnik D, Zahtila E (1991) The date shell *Lithophaga lithophaga* Linnaeus, and its habitat calls for protection in the Adriatic Sea. In: Boudouresque CF, Avon M, Gravez V (eds) *Les espèces marines à protéger en Méditerranée*. GIS Posidonie, Marseille, pp 151–158
- Hutchings PA (1986) Biological destruction of coral reefs. *Coral Reefs* 4:239–252
- Hutchings PA (2008) Role of polychaetes in bioerosion of coral substrates. In: Wisshak M, Tapanila L (eds) *Current developments in bioerosion*. Springer, Heidelberg, pp 249–264
- Jaccarini V, Bannister WH, Miccafe H (1968) The pallial glands and rock boring in *Lithophaga lithophaga* (Lamellibranchia, Mytilidae). *J Zool Lond* 154:397
- Kleemann KH (1973) Der Gesteinsabbau durch Ätzmuscheln an Kalkküsten. *Oecologia* 13:377–395
- Kroeker KJ, Micheli F, Gambi MC, Martz TR (2011) Divergent ecosystem responses within a benthic marine community to ocean acidification. *PNAS* 108:14515–14520
- Labura Ž, Hrs-Brenko M (1990) Infestation of European flat oyster (*Ostrea edulis*) by polychaete (*Polydora hoplura*) in the northern Adriatic Sea. *Acta Adriatica* 31:173–181
- Laubier L (1966) Le coralligène des Albères: monographie biocénotique. *Ann Inst Océanogr Monaco* 72:1–112
- Lauckner G (1983) Diseases caused by metazoans. Agents: Porifera. In: Kinne O (ed) *Diseases of Mollusca*, vol II, Introduction, Bivalvia to Scaphopoda. Biologische Anstalt Helgoland, Hamburg, pp 617–626
- Le Bris S, Le Campion-Alsumard T, Romano J-C (1998) Caractéristiques du feutrage algal des récifs coralliens de Polynésie française soumis à différentes intensités de bioérosion. *Oceanol Acta* 21:695–708
- Le Campion-Alsumard T (1975) Etude expérimentale de la colonisation d'éclats de calcite par les Cyanophycées endolithes marines. *Cah Biol Mar* 16:177–185
- Le Campion-Alsumard T (1979) Les Cyanophycées endolithes marines. Systématique, ultrastructure, écologie et biodestruction. *Oceanol Acta* 2:143–156
- Le Campion-Alsumard T, Golubic S, Hutchings PA (1995) Microbial endoliths in skeletons of live and dead corals: *Porites lobata* (Moorea, French Polynesia). *Mar Ecol Prog Ser* 117:149–157
- Le Grand HM, Fabricius KE (2011) Relationship of internal macrobioeroder densities in living massive *Porites* to turbidity and chlorophyll on the Australian Great Barrier Reef. *Coral Reefs* 30:97–107
- Loya Y (1982) Life history strategies of boring bivalves in corals. In: *Proceedings of the 4th international coral reef symposium*, Marine Science Centre, University of the Philippines, Manila, Philippines, vol 2, pp 756
- Maldonado M (1992) Demosponges of the red coral bottoms from the Alboran Sea. *J Nat Hist* 26:1131–1161

- Martin D, Britayev TA (1998) Symbiotic polychaetes: review of known species. In: Ansell AD, Gibson RN, Barnes M (eds) *Oceanogr Mar Biol Ann Rev* 36:217–340
- Mastrototaro F, D'Onghia G, Corriero G, Matarrese A, Maiorano P, Panetta P, Gherardi M, Longo C, Rosso A, Scutio F, Sanfilippo R, Gravili C, Boero F, Taviani M, Tursi A (2010) Biodiversity of the white coral bank of Cape Santa Maria di Leuca (Mediterranean Sea): an update. *Deep-Sea Res II* 57:412–430
- McLoughlin N, Furnes H, Banerjee NR, Staudigel H, Muehlenbachs K, de Wit M, Van Kranendonk MJ (2008) Micro-bioerosion in volcanic glass: extending the ichnofossil record to Archaean basaltic crust. In: Wisshak M, Tapanila L (eds) *Current developments in bioerosion*. Springer, Heidelberg, pp 371–396
- Melone N (1965) I poriferi associate a *Corallium rubrum* (L.) della Sardegna. *Ann Mus Civ Storia Nat 'Giacomo Doria' Genova* 75:343–358
- Micael J, Alves MJ, Costa AC, Jones MB (2009) Exploitation and conservation of echinoderms. *Oceanogr Mar Biol Ann Rev* 47:191–208
- Mitchell-Tapping HJ (1983) Experimental fracturing of various corals in the reef environment. *Caribb J Sci* 19:49–52
- Mokady O, Lazar B, Loya Y (1996) Echinoid bioerosion as a major structuring force of Red Sea coral reefs. *Biol Bull* 190:367–372
- Nava H, Carballo JL (2008) Chemical and mechanical bioerosion of boring sponges from Mexican Pacific coral reefs. *J Exp Biol* 211:2827–2831
- Naylor LA, Viles HA (2002) A new technique for evaluating short-term rates of coastal bioerosion and bioprotection. *Geomorphology* 47:31–44
- Neumann AC (1966) Observations on coastal erosion in Bermuda and measurements of the boring rate of the sponge, *Cliona lampa*. *Limnol Oceanogr* 11:92–108
- Palmer M, Fornós JJ, Balaguer P, Gómez-Pujol LI, Pons GX, Villanueva G (2003) Spatial and seasonal variability of the macro-invertebrate community of a rocky coast in Mallorca (Balearic Islands): implications for bioerosion. *Hydrobiologia* 501:13–21
- Perry CT, Macdonald IA (2002) Impacts of light penetration on the bathymetry of reef microboring communities: implications for the development of microendolithic trace assemblages. *Palaeogeogr Palaeoclimatol Palaeoecol* 186:101–113
- Pomponi SA (1980) Cytological mechanisms of calcium carbonate excavation by boring sponges. *Int Rev Cytol* 65:301–319
- Porras R, Bataller JV, Murgui E, Torregrosa MT (1996) Trophic structure and community composition of polychaetes inhabiting some *Sabellaria alveolata* (L.) reefs along the Valencia Gulf Coast, Western Mediterranean. *Mar Ecol* 17:583–602
- Privitera D, Chiantore M, Mangialajo L, Glavic N, Kozul W, Cattaneo-Vietti R, Giancarlo A (2005) Spatial scale of variability of sea urchins and algal communities in the Dubrovnik area. In: *Proceedings of the 15th meeting of the Italian Society of Ecology*, pp 1–6
- Pronzato R (2000) Il corallo. L'oro rosso del Mediterraneo. *Boll Mus Inst Biol Univ Genova* 64–65:1–94
- Pyko IA (2009) Ichnodiversity and bioerosion patterns in the Eastern Mediterranean – an experimental study. Masters, Friedrich-Alexander-Universität Erlangen-Nürnberg, 68 pp
- Ricci S, Priori GF, Bartolini M (2008) Bioerosione di pavimentazioni musive sommerse ad opera della spugna endolitica *Cliona celata*. *Boll Inst Sup Conserv Restauro* 15:7–18
- Risk MJ, MacGeachy JK (1978) Aspects of bioerosion of modern Caribbean reefs. *Rev Biol Trop suppl* 26:85–105
- Rodolfo-Metalpa R, Houlbrèque F, Tambutté É, Boisson F, Baggini C, Patti FP, Jeffree R, Fine M, Foggo A, Gattuso J-P, Hall-Spencer JM (2011) Coral and mollusc resistance to ocean acidification adversely affected by warming. *Nat Clim Chang* 1:308–312
- Rose CS, Risk MJ (1985) Increase in *Cliona delitrix* infestation of *Montastrea cavernosa* heads on an organically polluted portion of the Grand Cayman fringing reef. *PSZNI Mar Ecol* 6:345–363
- Rosell D, Uriz M-J, Martin D (1999) Infestation by excavating sponges on the oyster (*Ostrea edulis*) populations of the Blanes littoral zone (north-western Mediterranean). *J Mar Biol Assoc UK* 79:409–413
- Russo GF, Cicogna F (1991) The date mussel (*Lithophaga lithophaga*), a 'case' in the Gulf of Naples. In: Boudouresque CF, Avon M, Grave V (eds) *Les espèces marines à protéger en Méditerranée*. GIS Posidonie, Marseille, pp 141–150
- Rützler K, Rieger G (1973) Sponge burrowing: fine structure of *Cliona lampa* penetrating calcareous substrate. *Mar Biol* 21:144–162
- Santangelo G, Carletti E, Maggi E, Bramanti L (2003) Reproduction and population sexual structure of the overexploited Mediterranean red coral *Corallium rubrum*. *Mar Ecol Prog Ser* 248:99–108
- Santos A, Mayoral E, da Silva CM, Cachao M, Domènech R, Martinell J (2008) Trace fossil assemblages on Miocene rocky shores of southern Iberia. In: Wisshak M, Tapanila L (eds) *Current developments in bioerosion*. Springer, Heidelberg, pp 431–450
- Sartoretto S (1998) Bioérosion des concrétions coralligènes de Méditerranée par les organismes perforants: essai de quantification des processus. *CR Acad Sci Paris Sci Terre Planète* 327:839–844
- Sartoretto S, Francour P (1997) Quantification of bioerosion by *Sphaerichinus granularis* on 'coralligene' concretions of the western Mediterranean. *J Mar Biol Assoc UK* 77:565–568
- Schneider J (1976) Biological and inorganic factors in the destruction of limestone coasts. *Contrib Sed Geol* 6:1–123
- Schneider J (1977) Carbonate construction and decomposition by epilithic and endolithic micro-organisms in salt- and freshwater. In: Flügel E (ed) *Fossil algae*. Springer, Berlin, pp 248–260
- Schönberg CHL (2008) A history of sponge erosion: from past myths and hypotheses to recent approaches. In: Wisshak M, Tapanila L (eds) *Current developments in bioerosion*. Springer, Heidelberg, pp 165–211
- Schönberg CHL, Suwa R (2007) Why bioeroding sponges may be better hosts for symbiotic dinoflagellates than many corals. In: Custódio MR, Lôbo-Hajdu G, Hajdu E, Muricy G (eds) *Porifera research: biodiversity, innovation and sustainability*. Museu Nacional, Rio de Janeiro, Brazil Série Livros, vol 28, 569–580
- Schönberg CHL, Tapanila L (2006) Bioerosion research before and after 1996 – a discussion of what has changed since the first international bioerosion workshop. *Ichnos* 13:99–102
- Schönberg CHL, Wilkinson CR (2001) Induced colonization of corals by a clionid bioeroding sponge. *Coral Reefs* 20:69–76
- Schönberg CHL, Wisshak M (2012) The perks of being endolithic. *Aquatic Biol* 17:1–5
- Schönberg CHL, Suwa R, Hidaka M (2008) Sponge and coral zooxanthellae in heat and light: preliminary results of photochemical efficiency monitored with pulse amplitude modulated fluorometry. *Mar Ecol* 29:247–258
- Scott PJB (1991) Rapid destruction of PVC piping by boring bivalves. *Int Biodeterior* 27:87–92
- Scott PJB, Risk MJ (1988) The effect of *Lithophaga* (Bivalvia: Mytilidae) bore holes on the strength of the coral *Porites lobata*. *Coral Reefs* 7:145–151
- Scott PJB, Moser KA, Risk MJ (1988) Bioerosion of concrete and limestone by marine organisms: a 13 year experiment from Jamaica. *Mar Poll Bull* 19:219–222
- Simboura N, Zenetos A (2005) Increasing polychaete diversity as a consequence of increasing research effort in Greek waters: new records and exotic species. *Mediterr Mar Sci* 6:75–88
- Smyth MJ (1989) Bioerosion of gastropod shells: with emphasis of effects of coralline algal cover and shell microstructure. *Coral Reefs* 8:119–125
- Szmant AM (2002) Nutrient enrichment on coral reefs: is it a major cause of coral reef decline? *Estuaries* 25:743–766
- Tapanila L (2008) The endolithic guild: an ecological framework for residential cavities in hard substrates. In: Wisshak M, Tapanila L (eds) *Current developments in bioerosion*. Springer, Heidelberg, pp 3–20

- Thomson Reuters (2011) Web of knowledge. http://apps.webofknowledge.com/UA_GeneralSearch_input.do?product=UA&search_mode=GeneralSearch&SID=Q18Ilo6BF4i@pOobL6e&preferencesSaved=. Accessed 14 Sept 2011
- Tribollet A (2008a) The boring microflora in modern coral reef ecosystems: a review of its roles. In: Wisshak M, Tapanila L (eds) Current developments in bioerosion. Springer, Heidelberg, pp 67–94
- Tribollet A (2008b) Dissolution of dead corals by euendolithic organisms across the northern Great Barrier Reef (Australia). *Microbiol Ecol* 55:569–580
- Tribollet A, Golubic S (2005) Cross-shelf differences in the pattern and pace of bioerosion of experimental carbonate substrates exposed for 3 years on the northern Great Barrier Reef, Australia. *Coral Reefs* 24:422–434
- Tribollet A, Atkinson MJ, Langdon C (2006) Effects of elevated $p\text{CO}_2$ on epilithic and endolithic metabolism of reef carbonates. *Glob Chang Biol* 12:220–2208
- Tribollet A, Godinot C, Atkinson M, Langdon C (2009) Effects of elevated $p\text{CO}_2$ on dissolution of coral carbonates by microbial euendoliths. *Global Biogeochem Cycles* 23:1–7
- Tunncliffe V (1979) The role of boring sponges in coral fracture. In: Lévi C, Boury-Esnault N (eds) *Biologie des spongiaires*, vol 291, Colloques internationaux du centre national de la recherche scientifique. CNRS, Paris, pp 309–315
- Vénec-Peyré M-T (1996) Bioeroding foraminifera: a review. *Mar Micropaleontol* 28:19–30
- Vicente VP (1990) Response of sponges with autotrophic endosymbionts during the coral-bleaching episode in Puerto Rico. *Coral Reefs* 8:199–202
- Vogel J, Brett CE (2009) Record of microendoliths in different facies of the Upper Ordovician in the Cincinnati Arch region USA: the early history of light-related microendolithic zonation. *Palaeogeogr Palaeoclimatol Palaeoecol* 281:1–24
- Ward P, Risk MJ (1977) Boring pattern of the sponge *Cliona vermifera* in the coral *Montastrea annularis*. *J Paleontol* 51:520–526
- Warne JE (1975) Borings as trace fossils, and the process of marine bioerosion. In: Frey RW (ed) *The study of trace fossils*. Springer, Berlin, pp 181–229
- Weinberg S (1991) Faut-il protéger les gorgones de Méditerranée? In: Boudouresque CF, Avon M, Gravez V (eds) *Les espèces marines à protéger en Méditerranée*. GIS Posidonie, Marseille, pp 47–52
- Wilson MA (2006) Macroborings and the evolution of marine bioerosion. In: Miller W III (ed) *Trace fossils: concepts, problems, prospects*. Elsevier, Amsterdam, pp 356–367
- Wisshak M (2008) Two new dwarf *Entobia* ichnospecies in a diverse aphotic ichnocoenosis (Pleistocene/Rhodes, Greece). In: Wisshak M, Tapanila L (eds) *Current developments in bioerosion*. Springer, Heidelberg, pp 213–233
- Wisshak M (2012) Microbioerosion. In: Knaust D, Bromley RG (eds) *Trace fossils as indicators of sedimentary environments. Developments in sedimentology* 64. Elsevier, Amsterdam, pp 213–243
- Wisshak M, Tribollet A, Golubic S, Jakobsen J, Freiwald A (2011) Temperate bioerosion: Ichno- and biodiversity from intertidal to bathyal depths (Azores). *Geobiology* 9:492–520
- Wisshak M, Schönberg CHL, Form A, Freiwald A (2012) Ocean acidification accelerates reef bioerosion. *PLoS ONE* 7(9):e45124
- Xavier JR, Rachello-Dolmen PG, Parra-Velandia F, Schönberg CHL, Breeuwer JAJ, van Soest RWM (2010) Molecular evidence of cryptic speciation in the ‘cosmopolitan’ excavating sponge *Cliona celata* (Porifera, Clionaidae). *Mol Phylogenet Evol* 56:13–20
- Yonge CM (1963) Rock-boring organisms. In: Soggnæs RF (ed) *Mechanisms of hard tissue destruction*. Publ Am Association Advancem Sci 75: 1–24
- Zenetos A, Katsanevakis S, Poursanidis D, Crocetta F, Damalas D, Apostolopoulos C (2011) Marine alien species in Greek Seas: additions and amendments by 2010. *Medit Mar Sci* 12:95–120
- Zundevich A, Lazar B, Ilan M (2007) Chemical versus mechanical bioerosion of coral reefs by boring sponges – lessons from *Pione cf. vastifica*. *J Exp Biol* 210:91–96